

Large scale fluctuations in the distribution of galaxies from the Two degree Field Galaxy Redshift Survey

Francesco Sylos Labini^{1,2}, Nikolay L. Vasilyev³, Yuriy V. Baryshev³

¹ Centro Studi e Ricerche Enrico Fermi, Via Panisperna 89 A, Compendio del Viminale, 00184 Rome, Italy

² Istituto dei Sistemi Complessi CNR, Via dei Taurini 19, 00185 Rome, Italy.

³ Institute of Astronomy, St.Petersburg State University, Staryj Peterhoff, 198504, St.Petersburg, Russia

Received / Accepted

Abstract. We study statistical properties of galaxy structures in several samples extracted from the Two degree Field Galaxy Redshift Survey. In particular, we measure conditional fluctuations by means of the scale-length method and we determine their probability distribution. In this way we find that galaxy distribution in these samples is characterized by large amplitude fluctuations of large spatial extension, whose size is limited only by samples boundaries. These fluctuations are quite typical and persistent in the samples volumes, and they are detected in two independent regions in the northern and southern galactic caps. We discuss the relation of the scale-length method to several statistical quantities, such as counts of galaxies as a function of redshift and apparent magnitude. We confirm previous results which have determined, by magnitude and redshift counts, that there are fluctuations of the order of 30% between the southern and the northern galactic caps and we relate explicitly these counts to structures in redshift space. We show that the estimation of fluctuations amplitude normalized to the sample density is biased by systematic effects, which we discuss in detail. We consider the type of fluctuations predicted by standard cosmological models of structure formation in the linear regime and, to study non-linear clustering, we analyze several samples of mock galaxy catalogs generated from the distribution of dark matter in cosmological N-body simulations. In this way we conclude that the galaxy fluctuations present in these samples are too large in amplitude and too extended in space to be compatible with the predictions of standard models of structure formation.

Key words. Cosmology: observations; large-scale structure of Universe;

1. Introduction

In one of his seminal papers de Vaucouleurs (1970) put into an historical perspective the problem of galaxy large scale structures and the question about the scale where galaxy distribution turns to homogeneity¹. He pointed out that observations have firstly found that galaxies are not randomly distributed, then that in the fifties the same property was assigned to cluster centers and finally that at the end of the sixties the discovery of super-clusters has still enlarged the scale of structures in the universe thus pushing to larger and larger scales the scale where the approach to homogeneity occurs.

In the last twenty years many observations have been dedicated to the study of the large scale distributions of galaxies (Huchra et al., 1983; Falco et al., 1999; Giovanelli

and Haynes, 1993; da Costa et al., 1988; Shectman et al., 1996; York et al., 2000; Colless et al., 2001). Despite the fact that large scale galaxy structures, of size of the order of several hundreds of Mpc/h², have been observed (de Lapparent Huchra and Geller, 1986; Geller and Huchra, 1989; Giovanelli and Haynes, 1993; Gott et al., 2005) to be the typical feature of the distribution of visible matter in the local universe, the statistical analysis measuring their properties has identified a characteristic scale which has only slightly changed since its discovery fourty years ago in angular catalogs. This scale, r_0 , was measured to be the one at which fluctuations in the galaxy density field are about twice the value of the sample density and it was indeed determined to be $r_0 \approx 5$ Mpc/h in the Shane and Wirtanen angular catalog (Totsuji and Kihara, 1969). Subsequent measurements of this scale — see e.g. Davis and Peebles (1983); Davis et al. (1988); Park et al. (1994); Benoist et al. (1996); Norberg et al. (2001, 2002b); Zehavi

¹ Note that de Vaucouleurs simply considered the scale where galaxies approaches to a random distribution. A distribution can be homogeneous, i.e. with a well-defined mean density, without being random, i.e. there can be small amplitude fluctuations with weak long-range correlations.

² We use $H_0 = 100h$ km/sec/Mpc for the value of the Hubble constant.

et al. (2002, 2004) — found a similar value, although in several samples larger values of r_0 have been found (i.e. $r_0 \approx 6 - 12$ Mpc/h). This variation was then ascribed to a luminosity dependent effect — see e.g. Davis et al. (1988); Park et al. (1994); Benoist et al. (1996); Zehavi et al. (2002).

Theoretical models of galaxy formation, like the Cold Dark Matter (CDM) one (see Peacock, 1999) are able to predict the scale r_0 once it is given the amplitude and correlation properties of fluctuations of the initial conditions in the early universe. The normalization of the matter initial condition can be obtained by measuring the amplitude and correlation properties of the anisotropies of the Cosmic Microwave Background Radiation (CMBR). Then by calculating the evolution of small density fluctuations in the linear perturbation analysis of a self-gravitating fluid in an expanding universe, it is possible to predict the scale r_0 today. This turns out, in current models as the CDM ones, to be $r_0 \approx 5$ Mpc/h (Springel et al., 2005). On scales $r < r_0$ models are unable to make precise predictions on the shape of the correlation function because gravitational clustering in the non-linear regime is difficult to be treated. Gravitational N-body simulations are then used to investigate structure formation in the non-linear phase. In addition, given that models predict, for $r > r_0$, a precise type of small amplitude fluctuations, it is possible to simply relate, by using the linear perturbation analysis mentioned above, the properties of fluctuations in the present matter density field to those in the initial conditions. It results that in a certain range of scales larger than r_0 , small amplitude fluctuations should have still positive correlations. Particularly, for $r_0 < r < r_c$, fluctuations have very small amplitude and weak positive correlations (see Sylos Labini and Vasilyev, 2008). At even larger scales $r > r_c$ (where this is estimated from CMBR measurements to be $r_c \approx 100$ Mpc/h) all models predict that the matter density field presents *anti-correlations* which tends to zero with a (negative) power law behavior of the type $-r^{-4}$ (Gabrielli et al., 2002; Sylos Labini and Vasilyev, 2008). This negative power-law tail corresponds in real-space to the linear dependence of the matter power-spectrum (PS) on the wave-number, i.e. $P(k) \sim k$. This former represents a behavior which can be interpreted as a consistency requirement for the properties of density fluctuations in Friedmann-Robertson-Walker models (Gabrielli et al., 2002). Because of the change of sign of the correlation function at r_c , this length-scale represents the cut-off in the size of weak amplitude structures in standard models. Thus, in the regime where fluctuations are small and have weak positive correlations, i.e. for $r_0 < r < r_c$, the present matter density field reflects the imprint of the initial conditions.

The fundamental test for current models of galaxy formation then concerns whether density fluctuations at large scales (i.e. $r > 10$ Mpc/h) have small amplitude or not. In addition another important question concerns the detection of anti-correlations on scales $r > r_c \approx 100$ Mpc/h (Sylos Labini and Vasilyev, 2008). The primary problem to

be considered in this respect concerns the statistical methods used to measure the amplitude of fluctuations and the range of correlations. There has been an intense debate in the last decade concerning this crucial point (Sylos Labini et al., 1998; Wu et al., 1999; Hogg et al., 2004; Gabrielli et al., 2005; Baryshev and Teerikorpi, 2005; Sylos Labini et al., 2008). The problem is indeed the following. Before one determines, in a given volume, the amplitude of fluctuations with respect to the sample density, one must have firstly tested that the former quantity is stable, i.e. that it does not depend on the sample size and/or it does not present large fluctuations in different samples containing the same type of objects. Indeed, in the case the distribution present structures and fluctuations at all scales in a given sample, i.e. it is inhomogeneous, the sample density is not a well-defined descriptor (Sylos Labini et al., 1998, 2008). In this situation all statistical quantities that are normalized to the sample density are affected by systematic effects. For this reason, prior to the characterization of fluctuations with respect to the sample density, a fundamental test consists in measuring conditional³ correlation properties (Gabrielli et al., 2005). It has been found that conditional statistical quantities, such as the conditional number of points in spheres, indeed show *scaling* properties at small scales $r < 20$ Mpc/h, e.g. the former grows as a function of distance slower than the volume (Sylos Labini et al., 1998; Hogg et al., 2004; Vasilyev et al., 2006; Sylos Labini et al., 2007, 2008). This result implies that unconditional quantities are affected by systematic finite-size effects and thus they do not give a reliable and meaningful estimation of correlations and amplitude of fluctuations. In this situation the length-scale r_0 can be an artifact of a statistical analysis which implicitly assumes that the sample average is a meaningful estimation of the asymptotic density, i.e. it assumes that the distribution is homogeneous and fluctuations have small amplitude well inside the sample volume.

While estimations of real-space correlation properties can be affected by finite-size effects, this is not the case when one counts galaxies as a function of redshift or apparent magnitude. In this case indeed one does not normalize statistical quantities to the sample average and large fluctuations have been found both in redshift (Kerscher et al., 1999; Chiaki et al., 2003) and angular surveys (see Picard, 1991; Frith et al., 2003). In particular in a CCD survey of bright galaxies within the Northern and Southern strips of the 2dF Galaxy Redshift Survey (2dFGRS) conclusive evidences were found that there are fluctuations of the order $\sim 30\%$ in galaxy counts as a function of apparent magnitude (Buswell et al., 2004). Further since in the angular region toward the Southern galactic cap (SGC) a deficiency, with respect to the Northern galactic cap (NGC), in the counts below magnitude ~ 17 (in the B filter) was found, persisting over the full area of the APM and APMBGC catalogs, this

³ In what follows we discuss in detail the difference between conditional and unconditional statistical properties.

would be an evidence that there is a large void of radius of about 150 Mpc/h implying that there is more excess large-scale power than detected in the 2dFGRS correlation function⁴ (Norberg et al., 2001, 2002b) or expected in the CDM models. It is indeed evident that, because of the difference in the counts amplitude, and thus in the sample density, between the NGC and the SGC samples, the estimation of the sample density is not stable. Thus the above mentioned problems on the normalization of fluctuations amplitude to the estimation of the sample density should be studied in great detail.

In this paper we use the 2dFGRS to study fluctuations in galaxy distribution at large scales, and to determine their statistical properties in redshift and magnitude space. Our aim is to employ statistical descriptors which do not, implicitly or explicitly, make use of the normalization of fluctuations to the sample average, thus avoiding the a-priori assumption of homogeneity inside a given sample. Thus we determine conditional statistical properties, expanding our previous findings in this same survey (Vasilyev et al., 2006), and we find that the puzzle of the coexistence of difference in densities in the NGC and SGC volumes on large spatial scales with a relatively small typical length scale of a few Mpc, can be understood as due to finite size effects in the estimation of the correlation function.

The paper is organized as follows. In Sec.2 we discuss the 2dFGRS data and the procedure used to construct the sub-samples used for the statistical analysis. The methods employed to characterize homogeneous and inhomogeneous distributions are briefly reviewed in Sec.3. In Sec.4 we present the main results of the analysis considering in detail the problems related to the estimation of fluctuations' amplitude normalized to the sample density. The properties of fluctuations of the matter density field predicted by theoretical models, in the linear regime, are briefly reviewed in Sec.5. Then, to study expected non-linear fluctuations in standard theoretical models, we analyze the properties of mock galaxy catalogs, generated from the dark matter density fields resulting from a cosmological N-body simulation — the Millennium Run by Springel et al. (2005). Finally in Sec.6 we discuss the results of the analysis of the 2dFGRS samples and we outline our main conclusions.

2. The 2dFGRS samples

The 2dFGRS (Colless et al., 2001) measured redshifts for more than 220,000 galaxies in two strips in the SGC and in the NGC. The median redshift is $z \simeq 0.1$. The apparent magnitude corrected for galactic extinction in the b_J filter is limited to be $14.0 < b_J < 19.45$. The selection of

VL sample	R_{min}	R_{max}	M_{min}	M_{max}	N_g
SGC400	100	400	-20.8	-19.0	29373
NGC400	100	400	-20.8	-19.0	23208
SGC550	150	550	-21.2	-19.8	26289
NGC550	150	550	-21.2	-19.8	18030

Table 1. Main properties of the obtained VL samples: R_{min} , R_{max} are the chosen limits for the metric distance; M_{min} , M_{max} are the corresponding limits in the absolute magnitude; N_g is the number of galaxies in the sample.

the samples used in the analysis is described in detail in Vasilyev et al. (2006). Here we briefly summarize the main points:

- To avoid the effect of the irregular edges of the survey we select two rectangular regions: in the SGC there is a slice of size $84^\circ \times 9^\circ$ limited by $-33^\circ < \delta < -24^\circ$, $-32^\circ < \alpha < 52^\circ$, while the NGC slice is smaller, i.e., $60^\circ \times 6^\circ$, with limits $-4^\circ < \delta < 2^\circ$, $150^\circ < \alpha < 210^\circ$ (coordinates are equatorial). The solid angle are $\Omega = 0.20, 0.11$ steradians for the SGC and the NGC slices.
- We select galaxies in the redshift interval $0.01 \leq z \leq 0.3$, with redshift quality parameter larger or equal to three, in order to get high quality redshifts (Hawkins et al., 2003).
- We do not use a correction for the redshift completeness mask and for the fiber collision effects. In fact, completeness varies mostly nearby the survey edges which are excluded in our sample. We assume that fiber collisions do not make a sensible change in the small scales correlation properties given that we set our lower cut-off to 0.5 Mpc/h which is larger than 0.1 Mpc/h used by Hawkins et al. (2003).
- The metric distance is standardly computed as in Zehavi et al. (2002):

$$R(z) = \frac{c}{H_0} \int_{\frac{1}{1+z}}^1 \frac{dy}{y \cdot (\Omega_M/y + \Omega_\Lambda \cdot y^2)^{1/2}}, \quad (1)$$

where we use the standard model parameters $\Omega_M = 0.3$ and $\Omega_\Lambda = 0.7$ and c is the light speed.

- The absolute magnitude is computed as in Zehavi et al. (2002):

$$M = b_J - 5 \cdot \log_{10} [R(z) \cdot (1+z)] - K(z) - 25, \quad (2)$$

where $K(z)$ is the K-correction term (Hogg et al., 2002)

- To calculate the K-correction $K(z)$ we use relations obtained by Madgwick et al. (2002) and we apply them as in Vasilyev et al. (2006).
- The volume limited (VL) samples are identified by two limits in the metric distance $R_{min} < R < R_{max}$ and two corresponding limits in the absolute magnitude M_{min} and M_{max} . In Tab.1 we report the properties of the VL samples.

It is known that the completeness mask for the survey includes the effects not only of redshift incomplete-

⁴ This statistics, as mentioned above, normalizes the amplitude of fluctuations to the estimation of the sample density. From it the length-scale $r_0 \simeq 6 - 8$ Mpc/h is derived. In what follows we discuss in details the problems related to the estimation of this statistical quantity.

VL sample	R_{min}	R_{max}	M_{min}	M_{max}	N_g
SGC400	100	400	-20.6	-19.2	22872
NGC400	100	400	-20.6	-19.2	18407
SGC550	150	550	-21.1	-20.0	18890
NGC550	150	550	-21.1	-20.0	13495

Table 2. As Tab.1 but for the case in which more conservative cuts in apparent magnitudes, i.e. $14.5 < b_J < 19.3$, have been used for the selection of the galaxies

ness (which varies both with field and with apparent magnitude) but also of the variations in the faint limiting magnitude across the survey regions. In order to test the robustness of our results, without correcting directly for the incompleteness using ad-hoc assumptions, we have restricted the sample to the apparent magnitude limits $14.5 < b_J < 19.3$ (see Hawkins et al., 2003). We report in Tab.2 the characteristics of the VL with these more conservative cuts.

As a self-consistent test we note below that the statistical analysis in the samples constructed with less conservative cuts agree with previous determinations for what concerns the galaxy counts as a function of apparent magnitude Norberg et al. (2002a); Buswell et al. (2004), the redshift distribution Ratcliffe et al. (1998); Buswell et al. (2004) and the standard two-point correlation function Norberg et al. (2001, 2002b).

3. Statistical methods

In this section we review the main properties of stationary stochastic point processes. These include both the ones which have a strictly positive ensemble average density and those which have it equal to zero. This discussion allows to clarify which are the useful statistical methods to be employed, in both cases, for the analysis of a *finite sample*. A more exhaustive treatment can be found in Gabrielli et al. (2005).

3.1. Volume average, ensemble average and self-averaging property

The problem of the statistical characterization of correlations and fluctuations of a stochastic distribution of points in a finite sample of volume V , can be rephrased as the problem of measuring volume averaged statistical quantities. The basic issue concerns whether these are meaningful descriptors, i.e. whether they give or not stable statistical estimations of ensemble averaged quantities. In this respect one has to consider various problems which maybe clarified after the definition of the general probabilistic properties of stochastic point processes.

First of all we need to define the ensemble properties. In general it is assumed that galaxy distribution is a stationary stochastic process, which means that it is statistically translationally and rotationally invariant, i.e. it satisfies the condition of spatial statistical isotropy and ho-

mogeneity in order to avoid special points or directions⁵. Stationary stochastic distributions satisfy these conditions also when they have zero average density in the infinite volume limit (Gabrielli et al., 2005).

Due to the assumption of ergodicity (i.e. that the ensemble average is equal to the infinite volume average), the existence of a well-defined average density implies that for a single realization of the mass distribution the following limit is well defined

$$\lim_{R \rightarrow \infty} \frac{1}{V(R; \mathbf{x}_0)} \int_{V(R, \mathbf{x}_0)} n(\mathbf{r}) d\mathbf{r} = n_0, \forall \mathbf{x}_0. \quad (3)$$

where $V(R, \mathbf{x}_0) \equiv 4\pi R^3/3$ is the volume of the sphere of radius R , centered on an arbitrary point \mathbf{x}_0 . The constant n_0 is strictly positive for homogeneous (or uniform⁶) distributions and it is zero for inhomogeneous ones (e.g. fractals). It is then clear that all statistical quantities which are normalized to n_0 are meaningless when this is equal to zero in the ensemble average or in the infinite volume limit.

Keeping these mathematical properties in mind we have to consider the situation occurring when in a finite sample, of size $\sim V^{1/3}$, the distribution is not homogeneous. In this case the estimator of the average mass density gives a large relative error with respect to the ensemble value and it is thus systematically biased. This implies that only statistical averages conditioned to the fact that the origin of coordinates is a point of the set are well-defined.

Clearly a distribution can be inhomogeneous up to a length scale λ_0 ⁷ and homogeneous for $r > \lambda_0$. Then for $r < \lambda_0$ the distribution is characterized by large fluctuations, and the average density is not a well defined quantity if it is estimated in samples of size $V^{1/3} < \lambda_0$. Instead for $r > \lambda_0$ density fluctuations are small and the sample density, when the sample volume is such that $V^{1/3} > \lambda_0$, converges to the asymptotic (or ensemble) value. The precise behavior of this convergence is determined by the (weak) two-point correlation properties of the distributions, and the convergence will be slower when correlations are long-ranged (Gabrielli et al., 2005).

3.2. Probability distribution of conditional fluctuations

In Sylos Labini et al. (2008) we have introduced the scale-length (SL) analysis. This consists in the determination of the number $N(r; R_i)$ of galaxies in spheres of radius r ,

⁵ Because of the hypothesis of statistical stationarity, statistical quantities generally depend only of the scalar distance between points.

⁶ Note that in order to avoid confusion sometimes it is denoted as uniform a stationary stochastic point process with a positive ensemble average density and non uniform when this is equal to zero. Both are statistically homogeneous stochastic processes (Gabrielli et al., 2005).

⁷ See below for a precise definition.

centered on the i^{th} galaxy⁸ whose distance from the origin is R_i . That is,

$$N(r; R_i) = \int_{V(r; R_i)} n(\mathbf{s}) d\mathbf{s} \quad (4)$$

where the integral is performed over the spherical volume $V(r; R_i)$ of radius r centered on the i^{th} galaxy at distance R_i from the origin and $n(\mathbf{s})$ is the microscopic number density, which is a function of the space position \mathbf{s} .

When Eq.4 is averaged over the whole sample it gives an estimate of the average conditional number of galaxies in spheres of radius r

$$\overline{N(r)} = \frac{1}{M(r)} \sum_{i=1}^{M(r)} N(r; R_i), \quad (5)$$

where the sum is extended to the $M(r)$ galaxies, the i^{th} at radial distance R_i from us, whose separation from the boundaries of the sample is smaller or equal to r . In this way when r grows the number of values $M(r)$ over which the mean in Eq.5 is performed decreases with r , because only those galaxies for which the sphere is fully included in the sample volume are considered as centers. This estimator, known as the full-shell estimator (Sylos Labini et al., 1998; Kerscher, 1999; Gabrielli et al., 2005), has the advantage to make the weakest a-priori assumptions about the properties of the distribution outside the sample volume. Indeed one may use incomplete spheres, by counting the number of galaxies inside a portion of a sphere and weighting this for the corresponding volume of the spherical portion (Kerscher, 1999). However this method implicitly uses the assumption that what is inside the incomplete sphere is a statistically meaningful estimate of the distribution in the whole spherical volume. This is incorrect when the distribution presents large fluctuations. For example in the part of a spherical volume which lies outside the sample boundaries there can be a void or a large scale structure: in this situation the weighted estimation is biased (Gabrielli et al., 2005). When the full-shell estimator is used one should consider that there is an intrinsic selection effect related to the geometry of the samples, which are small portions of spheres: when r is large only the more distant part of the sample is explored by the volume average (Gabrielli et al., 2005; Sylos Labini et al., 2008). Indeed, for large sphere radii $M(r)$ decreases and the location of the galaxies contributing to the average in Eq.5 is mostly placed at radial distance in the range $\sim [R_{\min} + r, R_{\max} - r]$ from the radial boundaries of the sample at $[R_{\min}, R_{\max}]$. In addition, given the geometry of the samples, for large r galaxies contributing to $M(r)$ will also lie toward the center of the spherical portion.

When Eq.5 scales as

$$\overline{N(r)} \sim r^D \quad (6)$$

and $D = 3$ the distribution is homogeneous, while for $D < 3$ it has long-range power-law correlations (Gabrielli

et al., 2005). The scaling of Eq.6 with $D < 3$ can be interpreted as the signature that the distribution is a fractal. However there are point distributions which, by construction, are not fractal objects but which may exhibit a scaling of the type given by Eq.6 (see Gabrielli et al., 2005).

From Eq.6 we obtain that, in general, for $D < 3$ the conditional density scales as

$$\overline{n(r)} = \frac{\overline{N(r)}}{V(r)} \sim r^{D-3};$$

in this situation the average density is not a well defined quantity and the sample density depends on the sample size i.e. it does not give a meaningful estimation of the ensemble average density.

When a distribution is fractal (or generally inhomogeneous) at small scales and homogeneous at large scales then we can identify the homogeneity scale λ_0 to be the scale such that (Gabrielli et al., 2005)

$$\overline{N(r)} \sim r^3, \quad \forall r > \lambda_0. \quad (7)$$

Depending on the details of the crossover from the strongly correlated regime at $r < \lambda_0$ to the weakly correlated one at $r > \lambda_0$ different definitions of λ_0 can be adopted but they all satisfy the condition given by Eq.7. For instance one may define such a scale to be the one at which the rms fluctuation have twice the value of the average density. Note that if strong clustering occurs only for $r < \lambda_0 < V^{1/3}$, in this range of scales it is required to make the statistical analysis suitable to describe large fluctuations. Instead for $\lambda_0 < r < V^{1/3}$ usual unconditional quantities are well defined and their convergence to the ensemble values can be studied.

The estimator defined by Eq.5 gives the first moment (i.e. the average number of points in spheres of radius r) of the PDF of conditional fluctuations $f(N; r)$ computed, at fixed r , from the values $\{N(r; R_i)\}_{i=1, \dots, M(r)}$. Note that this is generally different from the PDF of unconditional fluctuations — considered by e.g., Saslaw (2000) — both for homogeneous and inhomogeneous distributions, the difference being more important in the former case.

Note that when a distribution becomes homogeneous, i.e. Eq.7 is satisfied, the PDF is expected to converge, in a finite volume, to a Gaussian function⁹ (Gabrielli et al., 2005), i.e.

$$f(N; r \gg \lambda_0) \simeq \frac{1}{\sqrt{2\pi\Sigma^2(r)}} \exp\left(-\frac{[N(r) - \overline{N(r)}]^2}{2\Sigma^2(r)}\right), \quad (8)$$

where $\overline{\Sigma^2(r)}$ is the estimation of the variance of the random variable $N(r)$ (see below).

⁹ This is clearly the case if the number of points is large enough, otherwise the PDF is described by the Poisson distribution

⁸ This is thus a conditional quantity.

The second moment of $f(N, V)$ gives the conditional variance. For inhomogeneous distributions this is such that

$$\overline{\delta(r)^2} \equiv \frac{\overline{\Sigma^2(r)}}{N(r)^2} = \frac{\overline{N(r)^2} - \overline{N(r)}^2}{N(r)^2} \sim 1, \quad (9)$$

where the last equality means that fluctuations are persistent (Gabrielli and Sylos Labini, 2001). On the other hand for homogeneous distributions, with any kind of small-amplitude correlations we have that (Gabrielli and Sylos Labini, 2001)

$$\overline{\delta(r)^2} \ll 1. \quad (10)$$

Note that when the sample density n_S is well defined, i.e. it does not sensibly depend on the sample volume, one may define the reduced two-point correlation function which can be written as (Gabrielli et al., 2005)

$$\xi(r) = \frac{1}{4\pi r^2} \frac{dN(r)}{dr} \frac{1}{n_S} - 1. \quad (11)$$

In this case the homogeneity scale can be, for instance defined to be the scale beyond which $\xi(r) < 1$, or the scale such that $\delta^2(\lambda_0) = 1$ (see Gabrielli et al., 2005).

4. Results from the 2dFGRS

In this section we present the results of the statistical analysis of the VL samples of the 2dFGRS catalog discussed in Sec.2. We start by presenting the SL analysis to then pass to the description of the determination of the PDF of conditional fluctuations. Furthermore we consider its first moment, i.e. the average number of points in spheres. To illustrate the usefulness of the SL analysis we consider its relation to the counts of galaxies as a function of radial distance and of apparent magnitude. This allows us to discuss in detail the relation between small scale two-point correlations and large scale properties of fluctuations in the galaxy density field. We then consider the finite-size effects which systematically affect the determination of fluctuations amplitude normalized to the sample density. In addition we compute the average of the SL determinations $\{N(r; R_i)\}_{i=1..M(r)}$ in bins of radial distance. The comparison of the behaviors in the NGC and SGC slices allows us to place a lower limit to the homogeneity scale.

4.1. The scale-length analysis

In Figs.1-4 it is shown the behavior of the SL analysis in the four 2dFGRS samples we considered. One may note that in all cases there are large density fluctuations in the correspondence of the location of galaxy large scale structures. In Fig.5 it is shown a three-dimensional plot of the same SL analysis reported in Fig.4. One may see how well structures are identified by this analysis.

The number of points $M(r)$ over which $N(r; R_i)$ is computed, as function of the sphere radius r is shown in Fig.6: when the sphere radius r gets larger, the $M(r)$




Fig. 1. From top to bottom the SL analysis for the sample NGC400 with $r = 5, 10$ Mpc/h.




Fig. 2. The same of Fig.1 but now for the sample NGC550 with $r = 10, 15$ Mpc/h.

decreases quite rapidly. This is due to the geometrical selection effect previously discussed. In addition note that solid angle of the SGC slice is twice that of the NGC slice, and thus for $r > 20$ Mpc/h, there are more center-points in the SGC samples than in the NGC ones.

Let us briefly discuss the main features that we detect in the various samples ¹⁰

- NGC400: there are large structures, which transversely cross the sample, at about 240 Mpc/h and at 260, 270, 290 and 320 Mpc/h. All have approximately the same thickness of about 30-40 Mpc/h. When the sphere radius is increased to $r = 10$ Mpc/h the most prominent

¹⁰ Note that, for instance, Eke et al. (2004); Einasto et al. (2006) have indeed identified with different methods, the same structures we observe




Fig. 5. The SL analysis for the SGC550 sample. On the X and Y axis it is reported the coordinate of the center of a sphere of radius $r = 10$ Mpc/h (centered on a galaxy) and on the Z axis the number of galaxies inside it. The mean thickness of this slice is about 50 Mpc/h. Large fluctuations in the density field traced by the SL analysis are located in the correspondence of large scale structures.




Fig. 3. The same of Fig.1 but now for the sample SGC400 with $r = 5, 10, 20$ Mpc/h.




Fig. 4. The same of Fig.1 but now for the sample SGC550 with $r = 10, 20, 30$ Mpc/h.

structure remains the one at about 250 Mpc/h, which

is not sampled anymore when $r = 15$ Mpc/h. This is




Fig. 6. Number of center-points $M(r)$ as a function of the sphere radius r in the various samples considered.

due to the fact that sphere centers are located toward the faraway boundaries of the sample, because of the geometrical selection effect previously discussed. For $r = 15$ Mpc/h, although only few points (i.e. $M \approx 3000$) are effectively considered as sphere centers, density fluctuations are still large, determining variation of a factor four in the determination of the number of points in spheres at different radial distances.

- NGC550: the structure at $R \approx 250$ Mpc/h is clearly visible even in this sample and other structures are present at larger radial distances $R > 350$ Mpc/h. Fluctuations are still large up to the largest sphere radius $r = 25$ Mpc/h where however the number of centers rapidly decreases for the geometrical selection effect discussed in Sec.3
- SGC400: for sphere radius $r = 10$ Mpc/h the situation is similar to the NGC400 case, except for the fact that the radial distances corresponding to the large variations in the density field are different. This shows that large scale structures and the corresponding large fluctuations detected by the SL analysis are quite typical of the galaxy distribution. There are two prominent large scale structures at radial distance of the order of $R \sim 320$ Mpc/h. Finally note that for $r = 20$ Mpc/h the sample is dominated by one of the two structures just mentioned, which corresponds to a variation of order five in $N(r; R)$.
- SGC550: A structure at $R \approx 320$ Mpc/h with thickness of about 40 Mpc/h is present inside this sample as well. In addition other structures are visible for $R > 400$ Mpc/h. For the largest sphere radius $r = 30$ Mpc/h, where only few points are considered as sphere centers, fluctuations in $N(r; R)$ are of order five and they are due to structures located at radial distances $R > 380$ Mpc/h.

In summary galaxy distribution in these samples is characterized by several large scale structures which cross

their volumes. These structures are typical and persistent, i.e. they are detected at different radial distances and in the two different directions toward the SGC and NGC. They correspond in the SL analysis to large fluctuations of the density field: the thickness of the structures is partially an intrinsic property and partially due to the 'Finger of God' effect — the redshift space extension of large clusters and super-clusters — convolved with the sphere size used to measure the conditional density. Typical velocity dispersions are up to $\sim 10^3$ km/s, although Eke et al. (2004) concluded that the typical velocity dispersion for 2dF groups and clusters is 260 km/s which corresponds only to 2.6 Mpc/h in additional radial extension. Thus while peculiar velocities can increase the size of radial structures, their effect cannot explain the whole range of distances which characterize the observed structures.

The maximum allowed sphere radius we considered, which, as discussed, is set by the geometry of the samples, is $r = 30$ Mpc/h. For this value of r we find fluctuations of order four in $N(r; R)$ and this allows us to conclude that the homogeneity scale is certainly larger than 40 Mpc/h. In what follows we reinforce this conclusion by considering suitable statistical measurements.

4.2. Probability distributions of conditional fluctuations

We now turn to the discussion of the PDF of conditional fluctuations. The behaviors in the various samples are shown Figs.7-10. Note that we limit the discussion to the case where the number of determinations $\{N(r, R_i)\}_{i=1, \dots, M(r)}$ at fixed sphere radius r is larger than few thousands points. For smaller numbers the measurement is affected by weak statistics and by finite-size effects, thus not giving to a statistically robust result. As discussed previously, when the sphere radius increases, there is a decrease of the number of centers, and thus a for large r , whose precise value is determined by the geometry of each of specific sample, the measurement is affected by large statistical and systematic effects (see Fig.6).

The first point to note is that in all cases the maximum of $f(N, r)$ is statistically stable, i.e. it does not change when it is computed in the whole sample or in two non-overlapping sub-samples with equal volume (each half of the sample volume) at small (sub-sample S_1) and at large (sub-sample S_2) radial distance (see Fig.7 and Fig.9).

The tail for large values of N is instead affected by the different fluctuations which are present in different sub-volumes. The trend is obvious: larger the fluctuations of $N(r; R)$ more extended toward large N values is the tail of $f(N, r)$. Note that in the deepest samples, e.g. SGC550, there is a single structure which dominates the distribution. However this is placed in the middle of the sample and for this reason, apparently, there is no a systematic difference of the PDF when this is computed in the two half volumes of the sample, one nearby and one faraway. However it is clear that in this situation the shape of the




Fig. 7. Probability density function $f(N, r)$ of the values $\{N(r; R_i)\}_{i=1, \dots, M(r)}$ for NGC400 in the whole sample and the best fit with a Gaussian function. The different sphere radius r is reported in the y-axis. The behavior in two non-overlapping sub-samples (labeled as S_1 and S_2 is also reported — see text for details. Poisson error bars are shown as a reference.

PDF will be strongly affected by this fluctuation. Indeed, in each sub-sample, for the largest sphere radius r we find that $f(N, r)$ is systematically distorted with respect to smaller sphere radii. This is due to fact that the volume average cannot explore properly the full sample because of the geometrical selection effect which, as discussed, is present in the determination of $N(r; R)$. To properly determine the PDF at scales $r > 20$ Mpc/h larger samples are thus required.

Note that in all cases the PDF is systematically different from a Gaussian function, but the case of NGC550 for which there is the weakest statistics, and it is characterized by a long N tail which is directly related to the large scale structures present in these samples. In addition note that the PDF differs in different samples, especially for $r > 10$ Mpc/h. This implies that, because of the weak statistics and small volumes, a clean determination of the PDF is not possible. For this reason we limit our discussion in what follows to the first moment of the PDF, leaving the determination of the second moment to further samples where spatial volumes will be larger.

In Fig.11 we show the collapse plot of the PDF in the various samples and for the different sphere radius considered. A rough fit of the large N tail is given by

$$f(N, r) \sim N^{-3}.$$

However this results requires better samples to be confirmed.

4.3. Conditional average number of galaxies in spheres

In Fig.12 we show the determination of the whole-sample average conditional number of points in spheres, i.e. Eq.5.




Fig. 8. The same of Fig.7 for the sample NGC550 with $r = 10, 15$ Mpc/h. For $r = 15$ Mpc/h there are only 3765 determinations.




Fig. 9. The same of Fig.7 for the sample SGC400 with $r = 5, 10, 20$ Mpc/h.

Given that the PDF in all samples are statistically stable, i.e. they do not show important systematic differences in different sub-volumes, the full sample volume average provides a meaningful statistical quantity. These behaviors are the same as the ones found by Vasilyev et al. (2006) for the average conditional density. We find that in all samples

$$\overline{N(r)} = Br^D, \quad (12)$$

with $D = 2.2 \pm 0.1$ and where B is a constant. This result is statistically robust up to $r = 30$ Mpc/h although for the SGC550 sample the analysis can reach, with a weaker statistics, $r \sim 40$ Mpc/h. In all cases, however, the determinations for scales $r > 20$ Mpc/h are done on a relatively




Fig. 10. The same of Fig.7 for the sample SGC550 with $r = 10, 20$ Mpc/h.




Fig. 11. Collapse plot of the $f(N, r)$ in the various samples and for the different sphere radius considered. The normalization of the different samples has been performed in an arbitrary way and on the X-axis the has arbitrary units.

small number of points (see Fig.6) and thus they can be subject of systematic fluctuations. Note that the amplitude between the NGC400 and SGC400 samples (which have higher number density than that of the deeper samples) differs for about $\sim 30\%$ (see Fig.13). This is due to the different fluctuations which are present in the different volumes and it implies that the samples are not large enough to make a precise determination of the amplitude. This difference is reflected in the number counts as a function of apparent magnitude as we discuss below. In the NGC550 and SGC550 samples, although the samples volume is larger, the constant B in Eq.12 varies of about $\sim 20\%$ (see Fig.13).

Let us briefly discuss the determination of the constant pre-factor B in Eq.12 (Joyce and Sylos Labini, 2001; Gabrielli et al., 2005) and the normalization of the conditional average density in different VL samples. To this aim one should consider the joint conditional probability of finding a galaxy of luminosity L at distance r from another galaxy, i.e. the (ensemble) conditional average number of galaxies $\langle \nu(L, r) \rangle_p dr dL$ with luminosity in the range $[L, L + dL]$ and in the volume element dr at distance r from an observer located on a galaxy. We can make the greatly simplifying assumption, that

$$\langle \nu(\mathbf{y}, L) \rangle_p = \phi(L) \times \langle n(\mathbf{y}) \rangle_p, \quad (13)$$

where $\langle n(\mathbf{y}) \rangle_p$ is the average conditional density and $\phi(L)$ is the luminosity function such that $\phi(L)dL$ gives the probability that a randomly chosen galaxy has luminosity in the range $[L, L + dL]$. Although there is a clear evidence that there is a correlation between them, and for instance the brightest elliptical galaxies are found in the center of rich galaxy clusters, it has been tested that this is nevertheless a reasonable assumption in the galaxy catalogs available so far. To treat properly the case where the correlation between galaxy positions and luminosities is taken into account, one would need to use the multi-fractal formalism. This goes beyond the scope of the present manuscript (see Gabrielli et al., 2005). Note that an additional, much stronger, assumption often adopted in Eq.13 is that the space density is a constant, i.e. $\langle n(\mathbf{r}) \rangle = const.$: this assumption is for instance at the basis of the so-called standard minimum variance estimator (Davis & Huchra, 1982). It is clear that we avoid to make this further assumption as we want to test whether the space density is (or can be approximated by) a simple constant.

Using Eq.13 we may write the conditional average number of galaxies as a function of distance (in case $D = const.$)

$$\begin{aligned} \langle N(r; L_{VL}^1 < L < L_{VL}^2) \rangle_p & \\ &= \int_0^r \int_{L_{VL}^1}^{L_{VL}^2} \langle \nu(\mathbf{y}, L) \rangle_p dL d\mathbf{y} \\ &= B r^D \int_{L_{VL}^1}^{L_{VL}^2} \langle \phi(L) \rangle_p dL = B_{VL} r^D, \end{aligned} \quad (14)$$

where $N(r; L_{VL}^1 < L < L_{VL}^2)$ is the number of galaxies in a sphere of radius r and with intrinsic luminosity in the range $[L_{VL}^1, L_{VL}^2]$, and B_{VL} is the amplitude of the number counts in the VL sample with these limits in absolute luminosity. Because the luminosity function has an exponential cut-off at L^* , VL samples containing brighter galaxies show a smaller B_{VL} . By knowing the shape of the luminosity function it is simple to normalize the different B_{VL} in different VL samples (Joyce and Sylos Labini, 2001).

4.4. Average in radial bins

We can now use the data obtained by the SL method to investigate whether there is a convergence to homogeneity




Fig. 12. Average number of points in spheres of radius r around a galaxy. The difference amplitude in samples with different limits in absolute magnitude is simply ascribed by the effect of the luminosity function (see text). Error bars are estimated by the sample dispersion on the average value.




Fig. 13. The same of Fig.12 but divided by the best fit power-law behavior $r^{2.25}$. The variation of the amplitude B in Eq.12 is more clear in this representation. The determination for $r > 20$ Mpc/h are subject to systematic fluctuations, due to the limited volume and the weaker statistics.

at some large scales $r > 30 \div 40$ Mpc/h, i.e., the largest sphere radius allowed by the geometry of the samples. This is done as follows. We divide the whole range of radial distances in bins of thickness ΔR and we compute the average

$$\overline{N(r; R, \Delta R)} = \frac{1}{M_b} \sum_{R_j \in [R, \Delta R]}^{j=1, M_b} N(r; R_j), \quad (15)$$

where the sum is extended to the M_b determinations of $N(r; R)$ such that the radial distance is in the interval range $[R, R + \Delta R]$. It is clear that for this measurement we have to use a small r in order to avoid overlapping in




Fig. 14. Behavior of the . Eq.15 with $r = 5$ Mpc/h for NGC400 and SGC400 and $\Delta R = 50$ Mpc/h. The error is computed through Eq.16.

space between neighboring bins in radial distance. We the variance on the mean is

$$\overline{\Sigma(r; R, \Delta R)^2} = \sum_{R_j \in [R, \Delta R]}^{j=1, M_b} \frac{\left(N(r; R_j) - \overline{N(r; R, \Delta R)} \right)^2}{M_b(M_b - 1)}. \quad (16)$$

Note that the error analysis in Eq.16 assumes that the $N(r; R_j)$ are independent, but in fact they are correlated. The errors on these points are therefore substantially under-estimated. However if there would be a trend toward homogenization the error done by neglecting this correlation will be smaller and so the under-estimate of the error bars. Only for the case of a highly correlated distribution Eq.16 underestimates the error bars, which than represent a lower limit of the “true” error bars.

The quantity given by Eq.15 and its error (Eq.16) provide an estimation of the number of points in spheres of radius r averaged in bin of thickness ΔR . We expect that, if the distribution converges to uniformity at a scale λ_0 , then correspondingly $\overline{N(r; R, \Delta R > \lambda_0)}$ does not show large fluctuations as a function of R .

Results for the four samples are shown in Figs.14-15. One may note that for the largest radial bin chosen $\Delta R = 75$ Mpc/h there is no trend for homogenization, rather the measurements in bins centered at different R wildly scatters, i.e. their values are outside the statistical error bars given by Eq.16. This shows that large scale structures have an amplitude which is incompatible with homogeneity at scales smaller than $\lambda_0 = \Delta R = 75$ Mpc/h.

It is interesting to note that the large fluctuations between the NGC and SGC samples can not be generated by some redshift-dependent effect, such as the inclusion of galaxy evolution in the computation of absolute magnitudes. Indeed such a correction, which is expected to be anyway small given that the redshifts involved do not exceed 0.2, would affect in the same way both samples. Thus




Fig. 15. As Fig.14 but for NGC550 and SGC550 with $r = 10$ Mpc/h and $\Delta R = 50$ Mpc/h.

by comparing the estimation of the density in bins in the same range of radial distances, we can conclude that the fluctuations we have detected are intrinsic to the distribution of galaxies in these samples. A similar argument can be done for the effect of different cosmologies in the computation of the metric distance.

4.5. Radial counts in VL samples

A complementary way to study fluctuations at large scales in galaxy redshift surveys is represented by the determination of the radial counts i.e. the counts of galaxies as a function of the radial distance in VL samples (Gabrielli and Sylos Labini, 2001). In order to have a statistical estimator and to evaluate fluctuations we have divided the angular area of the samples into $N_f = 20$ non-overlapping sub-fields of equal solid angle. For each we compute the differential radial density $n_i(R; \Delta R)$ in bins of thickness $\Delta R = 10$ Mpc/h, where i^{th} labels the sub-field and R is, as usual, the radial distance. We then compute the average

$$\overline{n(R; \Delta R)} = \frac{1}{N_f} \sum_{i=1}^{N_f} n_i(R; \Delta R) \quad (17)$$

and the sample variance

$$\overline{\sigma_n^2(R; \Delta R)} = \frac{1}{N_f - 1} \sum_{i=1}^{N_f} (n_i(R; \Delta R) - \overline{n(R; \Delta R)})^2. \quad (18)$$

Results are shown in Figs.16-17. We note that in the NGC400 sample the structure at 250 Mpc/h is visible as a relatively large fluctuation of $\overline{n(R; \Delta R)}$ and a corresponding large error. This means that this structure covers partially the angular area of the survey. In the NGC550 sample the radial density is more flattened, although there is a large dispersion. In the SGC400 sample the structures at 160 Mpc/h and 320 Mpc/h are identified as local enhancements of $\overline{n(R; \Delta R)}$. The same occurs for the SGC550




Fig. 16. The average differential radial density $\overline{n(R; \Delta R)}$ (Eq.17) and its error (Eq.18) for the samples NGC400 and SGC400.




Fig. 17. The same of Fig.16 for SGC550 and NGC550.

case where the same two structures are visible. By comparing Figs.16-17 with Figs.1-4 one may note that the SL analysis is a much more powerful method than the simple counting as a function of radial distance in tracing large scale galaxy structures.

4.6. Redshift distribution in the magnitude limit sample

By studying the redshift distribution in the Durham/UKST Galaxy Redshift Survey there have been found fluctuations in the observed radial density function are of the order of 50% occurring on ~ 50 Mpc/h scales (Ratcliffe et al., 1998; Buswell et al., 2004). In a similar way in the 2dFGRS (Buswell et al., 2004) two clear holes in the galaxy distribution were detected in the




Fig. 18. *Upper panel:* Radial density in bins of thickness 10 Mpc/h in the NGC magnitude limited sample. The most prominent features identified by the $N(r; R)$ analysis are visible also by the simple counting. There is a large structure at ~ 240 Mpc/h. By comparing this figure with Figs.1-4 one may appreciate the usefulness of the SL method in tracing structures *Bottom panel:* The same for the SGC magnitude limited sample. Here there are large structures at ~ 180 Mpc/h and ~ 320 Mpc/h.

ranges $0.03 < z < 0.055$, with an under-density of $\sim 40\%$, and $0.06 < z < 0.1$ where the density deficiency is about $\sim 25\%$. These two under-densities, detected in particular in the the 2dFGRS southern galactic cap (SGC), are also clear features in the Durham/UKST survey. Given that the 2dFGRS SGC field is entirely contained within the areas of sky observed for the Durham/UKST survey the similarities in the redshift distributions are both evidences of the same features in the galaxy distribution (Buswell et al., 2004).

We can now compare the redshift distribution in the magnitude limit sample with the results obtained by the SL analysis. In Fig.18 we report the counting of galaxies as a function of the radial distance, in bins of thickness 10 Mpc/h, in the whole magnitude limited samples. It is interesting to compare these behaviors with Figs.1-4. The SL method clearly identifies the same structures, which are visible in Fig.18 as peaks of the radial distribution. However the SL method is able to quantify the amplitude of these fluctuations and, by applying the statistical analysis presented above, to determine how typical structures are.

4.7. Magnitude counts

In Fig.19 we report the differential counts of galaxies as a function of apparent magnitude in the SGC and NGC. The behaviors are similar to those found by Norberg et al. (2002a); Buswell et al. (2004): the former paper concluded that there is a conclusive evidence that counts in




Fig. 19. Differential counts of galaxies, in bins of $\Delta m = 0.25$, as a function of apparent magnitude in the SGC and NGC. A reference line corresponding to $N(m) \sim 10^{\alpha m}$ with $\alpha = 0.5$ is reported.

the SGC are down by 30% relative to the NGC counts. We find the same difference and, as discussed, we can directly relate it to the large scale structures which are present in both samples. Indeed, as already discussed, the amplitude of the conditional number of galaxies (i.e., Eq.12) is $\sim 20 \div 30\%$ higher for the NGC samples than for the SGC ones. In other words, in the NGC samples there are more structures, and thus fluctuations in the $N(r; R)$, than in SGC samples as can be seen by comparing, for instance, Fig.1 and Fig.3.

4.8. The two-point correlation function

The standard way to measure two-point correlations is accomplished by determining the function $\xi(r)$ given by Eq.11 — see e.g., Totsuji and Kihara (1969); Davis and Peebles (1983); Park et al. (1994); Benoist et al. (1996); Zehavi et al. (2002, 2004); Norberg et al. (2001, 2002b). As already mentioned when measuring this quantity it is implicitly assumed that the distribution is homogeneous well inside the sample volume, i.e. $\lambda_0 \ll V^{1/3}$. Let us see what happens when, inside a spherical sample of radius R_s there is a fractal distribution with dimension $D < 3$. We may estimate the sample density n_S (which is not an average quantity) by

$$n_S = \frac{N}{V} = \frac{3B}{4\pi} R_s^{D-3}, \quad (19)$$

where, in the second equality on the r.h.s., we used Eq.12 for the number of points in spheres. Eq.19 shows that the sample density depends on the sample size when $D < 3$. The estimator of the two-point correlation function can be written as (Gabrielli et al., 2005)

$$\overline{\xi(r)} + 1 = \frac{\overline{N(r, \Delta r)}}{V(r, \Delta r)} \cdot \frac{1}{n_S}. \quad (20)$$

The first ratio in the r.h.s. of Eq.20 is the average conditional density, i.e., the number of galaxies in shells of thickness Δr averaged over the whole sample, divided by the volume $V(r, \Delta r)$ of the shell. The second ratio in the r.h.s. of Eq.20 is the sample density. By using the second equality on the r.h.s. of Eq.19 for n_S and Eq.12 for the conditional number of points in spheres, we find

$$\overline{\xi(r)} = \frac{D}{3} \left(\frac{r}{R_s} \right)^{D-3} - 1, \quad (21)$$

which shows that the amplitude of $\xi(r)$ depends on R_s for $D < 3$, i.e. as long as the conditional density is a power-law as a function of scale ¹¹.

Note that Eq.21 has been obtained by making the assumption that the estimation of the sample density is given by the second equality in the r.h.s. of Eq.19. This is in general not the case, as the sample density in inhomogeneous distributions is subjected to fluctuations of order one at the scale of the sample size. Thus the behavior given by Eq.21 should be interpreted as giving a very rough estimation of the amplitude of $\overline{\xi(r)}$ in a finite sample when inside it there is a fractal distribution.

It is worth noticing that if the conditional density is a power-law function of scale than $\xi(r)$ does not have the same scale-dependent behavior. Indeed, as shown by Eq.21, $\overline{\xi(r)}$ has a break of the power law and it is possible to compute analytically the exponent of $\overline{\xi(r)}$ as a function of the exponent of the conditional density and the scale ratio r/R_s (Gabrielli et al., 2005). It is easy to show that at scales $r/R_s \ll 1$ the correlation exponent measured by the two-point correlation analysis coincides with that measured by the conditional density, while at larger scales the exponent measured by the $\xi(r)$ analysis is generally smaller than $D - 3$. This is indeed the result obtained by Hawkins et al. (2003) (see their Fig.6).

Let us now evaluate the sample density simply as $n_S = N/V$ in SGC400 and NGC400. Given that in our case the sample geometry is a sphere portion, the volume is given by

$$V_{2dFGRS} = \frac{\Omega}{3} (R_{max}^3 - R_{min}^3). \quad (22)$$

By using the parameters of the VL samples (see Tab.1) we obtain respectively in the SGC400 sample

$$\overline{n_{sgc}} = 7.0 \cdot 10^{-3}$$

galaxies per $(\text{Mpc}/h)^3$ and in the NGC400 sample

$$\overline{n_{ngc}} = 1.0 \cdot 10^{-2}.$$

Thus there is a $\sim 30\%$ difference in the amplitude. When we normalize the conditional density in the NGC400 and SGC400, which as already discussed have the same $\sim 30\%$

¹¹ Pair-counting based estimators, like the Davis & Peebles and/or the Landy & Szalay estimator, have the same pathologies when the conditional density is a power-law as a function of scale (Gabrielli et al., 2005).




Fig. 20. Standard two-point correlation function in the SGC400 and NGC400 sample estimated by Eq.20: the sample density is simply computed as N/V .

difference in amplitude, to the respective values of the sample density we get the amplitude of $\overline{\xi(r)}$ is nearly the same (see Fig.20). This is because both the nominator and denominator in the r.h.s. of Eq.20 vary in the same way, having a difference of the order of 30% in the two samples. Thus by measuring the amplitude of fluctuations normalized to the sample density the information about the large variations in the two different slices is lost. This *does not* however imply that fluctuations are small beyond $r_0 \approx 6$ Mpc/h as one would conclude by only considering the two-point correlation function. Actually, as we have discussed in the previous sections, fluctuations are large and persistent over the whole volume of these samples. It is indeed the normalization of fluctuations amplitude to a not well-defined quantity, the sample density, which in this case gives raise to the determination of the length scale $r_0 \approx 6$ Mpc/h, which is however a spurious quantity. Note that the volume of the SGC400 sample is larger than that of the NGC400 sample: this is reflected in the fact that the break of $\overline{\xi(r)}$ occurs at a larger distance than for the NGC400 sample.

For the samples NGC550 and SGC550 there is a similar situation but the variation of both the sample density and of the amplitude of the conditional density is of the order of the 20%.

Different estimators of the two-point correlation function, such the Davis and Peebles (DP) (Davis and Peebles, 1983) estimator and the Landy and Szalay (LS) (Landy and Szalay, 1993) estimator — see discussion in Sylos Labini and Vasilyev (2008) for more details about the different estimators —, give rise to an estimation of the amplitude of $\overline{\xi(r)}$ which agrees to the one just discussed above. For instance in Fig.21 it is shown a comparison of the different estimators. It is worth noticing that the different estimators give the sample amplitude of $\overline{\xi(r)}$ but they differ in the scale at which $\overline{\xi(r)}$ has the break in the




Fig. 21. Standard two-point correlation function in the SGC400 measured by means of different estimators, namely the Full-Shell (FS), the Davis and Peebles (DP) and the Landy and Szalay (LS).

power-law behavior. This is explained by the different way the estimators treat the boundary conditions, and, particularly, include (implicitly) the global condition known as integral constraint — see discussion in Sylos Labini and Vasilyev (2008).

The sample density n_S can be estimated differently from Eq.19. The only condition that it is required for Eq.20 to be a valid estimator is that the size r^* of the volume entering in the denominator of Eq.19 to be larger than the homogeneity scale λ_0 . If fact when $r^* \gg \lambda_0$ the estimation of the sample density will not differ substantially from its ensemble average value because, in this situation, the amplitude of the two-point correlation is, by definition, much smaller than unity. Thus we may consider another estimator of the two-point correlation function, which is the one introduced by Sylos Labini et al. (2008)

$$\overline{\xi(r; R, \Delta R)} + 1 = \frac{\overline{N(r, \Delta r)}}{V(r, \Delta r)} \cdot \frac{V(R, \Delta R)}{N(R, \Delta R)}, \quad (23)$$

where the second ratio on the r.h.s. is the density of points in a shell of thickness $r^* = \Delta R$ around the radial distance R . If the distribution is homogeneous, i.e., $r^* > \lambda_0$, and statistically stationary, Eq.23 should be statistically independent on the range of radial distances $(R, \Delta R)$ considered. In Fig.22 we show the determination of Eq.23 for different values of $R, \Delta R$ in the two deepest samples. One may note that the amplitude of $\overline{\xi(r)}$ systematically depends on the choice of the normalization and this is just the imprint of the large scale inhomogeneities present in these samples.

In summary, despite the fact that there is a relatively large difference in the densities between the NGC and the SGC, the amplitude of the correlation function is similar because it is measured with respect to a varying density, i.e. its value reflects the assumption of homogeneity which




Fig. 22. Standard two-point correlation function in the SGC550 and NGC550 samples estimated by Eq.23: the sample average density is computed in spheres of radius r^* and considering all center-points lying in a bin of thickness ΔR centered at different radial distance R .

is used in the definition in the ξ -analysis. Only by making an analysis of fluctuations not normalized the sample density one may detect the effect of the large spatial inhomogeneities characterizing these galaxy samples.

4.9. Analysis in the catalog with conservative cuts in apparent magnitude

We now briefly discuss the results of the SL analysis for the VL samples obtained with conservative magnitude cuts (see Tab.2). In Fig.23 we show the results of the SL analysis: by comparing this figure with Figs.1-4 by a simple visual inspection one may note that structures are extremely similar. Clearly, because of the lower number of points contained in the conservative-cuts samples, the value of $N(r; R)$ is different. The behavior of the PDF for the various samples is shown in Fig.23. The results are thus statistically stable.

5. Comparison with mock catalogs

Standard theories of galaxy formation assume that fluctuations of the matter density field in the early universe have very small amplitude, (Peebles, 1980; Peacock, 1999). In this situation, by denoting as $P(k, t)$ the PS of matter at an arbitrary time $t \gg 0$, where $t = t^*$ corresponds to some early times in the universe as we discuss below, the prediction of standard cosmological models can be written in this form

$$P(k, t) = AP(k, t^*)g(k, t). \quad (24)$$

Let us now discuss each of the three terms in the r.h.s. of Eq.24. It is possible to write $P(k, t^*) = P_{hz}(k)T^2(k)$ where $P_{hz}(k)$ is the Harrison-Zeldovich (HZ) PS and $T(k)$




Fig. 23. SL analysis for the samples with conservative apparent magnitude cuts. The value of the sphere radius is reported in the captions




Fig. 24. PDF of the samples with conservative apparent magnitude cuts. The value of the sphere radius is reported in the labels. In the comparison we used the normalized variable z , by using the transformation described by Eqs.26-27

is a transfer function which depends on the type of coupling between matter and radiation in the early universe (Peacock, 1999). The HZ PS is predicted by inflationary theories to be the outcome of the exponential amplification of quantum fluctuations occurring in the early phases after the Big-Bang and this is such that $P_{hz}(k) \propto k$. A PS behaving as $P(k) \propto k^n$ at small k with $n \geq 1$ belong to the class of super-homogeneous, or hyper-uniform (Torquato and Stillinger, 2003), distributions (Gabiellini et al., 2002). Their main characteristic is that density fluctuations are of surface type, i.e. that the relative mass variance shows the fastest possible decay as a function of scale $\sigma_{hz}^2(r) \propto 1/r^4$. The PS must have a cut-off at some large values of k , i.e.

at small scales. This is provided by the transfer function $T(k)$ which can be determined given the supposed properties of (dark) matter. For CDM this is such that at small k the PS maintains the HZ tail, while at large k it is approximately decays as k^{-2} (Peacock, 1999). Correspondingly the two-point correlation function is positive for $r < r_c$ and negative for $r > r_c$ with a power-law tail of the type $\xi(r) \sim -r^{-4}$ (Sylos Labini and Vasilyev, 2008).

Gravitational clustering in the linear regime, in an Einstein de Sitter cosmology, is characterized by a growing and a decaying mode, both of them power laws in time (Peebles, 1980) $g(k, t) = g(t) \propto t^{4/3}$. The amplitude A in Eq.24 is determined from the observations of CMBR anisotropies and from the theoretical assumptions on the nature of cosmological dark matter. From the time dependence of $P(k, t)$ it is possible to derive the time dependence of $\lambda_0^m(t)$, defined to be $\sigma^2(\lambda_0^m) = 1$. This grows as a power-law function of time as well: particularly for power-law PS, i.e. $P(k) \sim k^n$ and $n < 4$, one obtains (Peebles, 1980)

$$\lambda_0^m(t) \propto t^{\frac{3}{3+n}}. \quad (25)$$

To summarize the situation, we may identify at the present time two length-scales. The first is the homogeneity (or non-linearity) scale λ_0^m : for $r < \lambda_0^m$ non-linear clustering took place and thus it changes the shape of PS in a non-linear way. On the other hand for $r > \lambda_0^m$ fluctuations are still in the regime today and linear perturbation theory predicts a simple linear amplification of primordial correlations. From the normalization of the initial amplitude A to the CMBR anisotropies it results that $\lambda_0^m \approx 10$ Mpc/h (Springel et al., 2005; Spergel et al., 2007). In addition from the properties of the PS of CMBR anisotropies it is derived that the second intrinsic length scale of these models is about $r_c \approx 100$ Mpc/h (Sylos Labini and Vasilyev, 2008).

In order to study gravitational structure formation in the non-linear phase the common practice is to perform N-body simulations of theoretical models. This is done by integrating the equation of motions of N self-gravitating particles, in a volume V and by making use of periodic boundary conditions to represent an infinite (periodic) system. Initial particle correlations are given according to a given theoretical model, and the initial redshift is generally $z > 10$. The simulation is then run up to $z = 0$. In addition the space background is expanding and thus one follows the particles' motion in comoving coordinates. Particles are supposed to simulate the motion of fluid elements of the underlying dark matter field.

To identify galaxies one uses a phenomenological approach. As discussed, galaxies are supposed to form in the highest density peaks of the dark matter field. Thus when the simulation has reached the redshift $z = 0$ one uses semi-analytic models to identify galaxies. Among the largest simulations made public available, the Millennium run (Springel et al., 2005) used more than 10 billions particles to trace the evolution of the matter distribution in

a region of the universe in a cubic box of 500 Mpc/h. Semi-analytic catalogs constructed from the Millennium run contains about 10 million objects (Croton et al., 2006).

Here we analyze the semi-analytic catalog containing 9,925,229 objects in which the absolute magnitudes of mock galaxies are given in the BVRIK filters. To reproduce the same limits in absolute magnitude of the volume limited samples of the 2dFGRS we used the relations between magnitude in different filters given by Colless et al. (2001). In this way we have selected respectively 1,119,434 and 368,619 galaxies in a 500 Mpc/h cube. We then selected three slices with the same geometry of the real 2dFGRS samples which are hereafter called as SGC400m, NGC400m and NGC550m. The remaining sample, SGC550m, is 60° wide instead of 84° as in real data. The number of objects in each of the four samples is of the order of the one in the correspondent real 2dFGRS sample.

In Fig.25 we show the behavior of the SL analysis for the mock sample SGC550m in real and redshift space for $r = 20$ Mpc/h (see for comparison Fig.2 and Fig.4). One may note the effect of peculiar velocities is that of enhancing a little the structures which appear in real space. That the difference between real and redshift space is small for sphere radius $r \geq 10$ Mpc/h is shown in Fig.26 and Fig.27, where we plot the PDF of conditional fluctuations. In this case we have used the normalized variable

$$z_i(r) = \frac{N_i(r) - \overline{N}(r)}{\overline{\Sigma}(r)} \quad (26)$$

and we thus determine its PDF, that is

$$P(z_i, r) = f\left(N_i(r) = \overline{N}(r) + z_i \overline{\Sigma}(r)\right) \times \overline{\Sigma}(r) \quad (27)$$

where $f(N, r)$ is the PDF of the variable $N_i(r)$, $\overline{N}(r)$ its estimated first moment and $\overline{\Sigma}(r)$ its estimated standard deviation at the scale r . It is worth noticing that for sphere radius smaller than the homogeneity scale predicted by theoretical models, i.e. $\lambda_0^m = 10$ Mpc/h, the PDF has a relatively large tail for large values of N , although around the peak it is well fitted by Gaussian function. For $r > 10$ Mpc/h the PDF rapidly converges to a Gaussian and already for $r = 20$ Mpc/h the fit with a Gaussian function is extremely good also at large N . Correspondingly the plot of the SL analysis does not show the presence of large amplitude fluctuations.

In Fig.28 and Fig.29 we show the comparison between the PDF in the mock sample SGC500m (redshift space) and the one in the real sample SGC550. One may note that both for the sphere radius $r = 10$ Mpc/h and $r = 20$ Mpc/h, fluctuations are more persistent in real samples than in mock samples and this is reflected in the relatively fat tail of the PDF for large N values.

Fig.30 (upper panel) it is shown the behavior of the whole-sample average number of points in spheres, i.e. Eq.5. One may notice that for $r > \lambda_0^m = 10$ Mpc/h, the function $\overline{N}(r)$ grows with an exponent equal to the




Fig. 25. SL analysis with sphere radius $r = 20$ Mpc/, for the mock sample SGC550m (the samples with better statistics) in real space (*upper panel*) and in redshift space (*bottom panel*).




Fig. 26. PDF for the mock samples SGC400m and NGC400m in real space (RS) and redshift space (ZS) for $r = 10$ Mpc/h. The best fit with a Gaussian function is reported as reference.

space dimension (i.e., $D = 3$). At smaller scales the exponent is instead $D = 1.2 \pm 0.1$.

Finally in the Fig.30 (bottom panel) it is shown the behavior of of Eq.15 with $r = 5$ Mpc/h for the mock galaxy samples in real space. One may notice that fluctuations, when averaged over radial bin of thickness $\Delta R = 50$ Mpc/h are more smooth and quiet than the real samples and thus that there are no large fluctuations at scales larger than λ_0^m . This is clearly consistent with the behavior of the PDF previously described. The small difference for small R is due to the lower statistics at the nearby boundary of the samples.

We note that because the distribution becomes uniform at scales $r > \lambda_0 \approx 10$ Mpc/h the two-point correlation function is well-defined, both in real and redshift




Fig. 27. The same as Fig.26 but for the case $r = 20$ Mpc/h and for the mock samples SGC550m and NGC500m.




Fig. 28. Comparison between the PDF in the mock sample SGC400m (redshift space) and the one in the real sample SGC400. Poisson error bars are displayed as a reference.




Fig. 29. The same as Fig.28 but for the case $r = 20$ Mpc/h and for the sample SGC550.




Fig. 30. *Upper panel:* Whole-sample average number of points in spheres, i.e. Eq.5 for the mock samples in real space. *Bottom panel:* Behavior of Eq.15 for the mock galaxy samples in real space for SGC550m; the sphere radius is $r = 10$ Mpc/h and the radial bin thickness is $\Delta R = 75$ Mpc/h. The error is computed through Eq.16.

space. This presents an intrinsic length scale $\xi(r_0) = 1$ at about $r_0 \approx 6$ Mpc/h and beyond that scale it rapidly decays to zero.

The whole-sample average number of points in spheres (i.e. Eq.5) does show a systematic difference with respect to the real-space case. Indeed, while for $r > \lambda_0^m = 10$ Mpc/h it shows again a power-law behavior with $D = 3$, at smaller scales the exponent is $D = 2.2 \pm 0.1$ instead of $D = 1.2$ as for the real space case (see Fig.31 — upper panel). In addition the transition between the two regimes is faster than for the real space case.

Finally in Fig.31 (bottom panel) it is reported the behavior behavior of Eq.15 for the mock samples in redshift space. As for the real-space case, and consistent with the behaviors of the PDF in redshift space, fluctuations are substantially smoothed when averaged over radial bin of size $\Delta R > \lambda_0^m$.

6. Discussion and Conclusions

In this paper we have characterized fluctuations and correlations in the galaxy distribution, by considering several samples from the 2dFGRS (Colless et al., 2001). We have used statistical quantities which measure amplitude of fluctuations not normalizing it to the sample average. We have shown that estimations of the sample average are affected by systematic variations which are the imprint of the structures of large spatial extension present in these samples which correspond to large amplitude fluctuations. In particular we applied the scale-length (SL) method, introduced in Sylos Labini et al. (2008), to identify structures in redshift space. Then we have measured the probability density function (PDF) of conditional fluctua-




Fig. 31. *Upper panel:* Whole-sample average number of points in spheres, i.e. Eq.5. for the mock samples in redshift space. *Bottom panel:* Behavior of Eq.15 for the mock samples in redshift space for SGP550m; the sphere radius is $r = 10$ Mpc/h and the radial bin thickness is $\Delta R = 75$ Mpc/h. The error is computed through Eq.16.

tuations, at different scales r , and its first moment. We have found that the PDF is, in all considered samples and up to $r = 20$ Mpc/h (the largest distance scale we considered for this statistics), wildly different from a Gaussian function and characterized by a long tail, possibly with a power-law decay, signaling the presence of large fluctuations (e.g. structures) in the density field. We found that these fluctuations are persistent up to scales limited by sample sizes. By considering the first moment of the PDF we measured that galaxy distribution has a fractal dimension $D = 2.2 \pm 0.1$ up to $r \approx 40$ Mpc/h: at the largest distance scale we considered, i.e. for $r > 20$ Mpc/h, determinations are affected by relatively large fluctuations which reflect the intrinsic systematic error induced by structures, and thus the results have weaker statistical significance for the limited volume and number of points contributing to the average. This result is in agreement with our previous findings in Vasilyev et al. (2006). We have also placed a lower limit to the homogeneity scale, i.e. the scale beyond which fluctuations become small and the sample density estimation approaches the value of the (ensemble) average density, at $\lambda_0 = 70$ Mpc/h. This is because fluctuations on scales of the order of 70 Mpc/h are incompatible with a transition to homogeneity at that scale. This result is indeed completely consistent with the results by Buswell et al. (2004); Ratcliffe et al. (1998). The main results obtained in this survey are in very good agreement with a similar analysis of galaxy samples extracted from the SDSS (Sylos Labini et al., 2008).

We then compared the results obtained by the SL method with galaxy counts as a function of the radial distance and of the apparent magnitude. In this respect

we have discussed the fact that the SL method is very suitable an effective in tracing structures in redshift space and to measure the amplitude and distribution of density fluctuations. We have shown that galaxy counts results can be easily related to the results obtained by the SL method. Indeed we confirmed that the variation of about $\sim 30\%$ between galaxy counts as a function of apparent magnitude in the northern and southern angular regions of this survey, detected by Buswell et al. (2004), can be easily related to the different structures present in the different samples. Moreover such a large fluctuations make any estimation of the sample average, at scales comparable to the samples sizes, affected by important systematic effects.

In this respect we discussed the problems related to the estimation of the two-point correlation function, which normalizes the amplitude of fluctuations (as a function of scale) to the estimation of the sample average. We pointed out that apparently the amplitude of this function is not affected by fluctuations, and that this is stable in the Northern and Southern samples. However we have shown that this occurs because of the normalization to the sample average (i.e. the denominator of $\xi(r)$) which indeed shows the same large variation between the two angular regions as the conditional density (i.e. the nominator of $\xi(r)$). Thus we have concluded that the normalization of fluctuations amplitude to the sample average hides the presence of large differences between the two samples, being biased by uncontrolled systematic finite-size effects. In this situation the amplitude of the two-point correlation function does not give a reliable and meaningful estimation of the statistical properties of galaxy distribution in these samples: neither of the spatial extension of correlations nor of amplitude of fluctuations. It is indeed the normalization of fluctuations amplitude to a not well-defined quantity, the sample density, which in this case gives raise to the determination of the spurious length scale $r_0 \approx 6$ Mpc/h.

We then compared the results obtained by studying conditional fluctuations in the 2dFGRS samples with those obtained in mock galaxy catalogs. These are constructed from the dark matter density field in cosmological simulations which have been evolved to study the effect of non-linear gravitational clustering in an expanding universe. We used the mock galaxy samples provided by Croton et al. (2006) and obtained from the largest N-body simulations publicly available at the moment (Springel et al., 2005). We have studied the mock galaxy catalogs both in real and redshift space, so to take into account the distortions due to the effect of peculiar velocities. We found that the PDF of conditional fluctuations is weakly affected by peculiar velocities and that it rapidly converges to a Gaussian function at scales $r > 10$ Mpc/h, in agreement with predictions of theoretical models both in real and redshift space. At smaller scales, i.e. $r < 10$ Mpc/h, the PDF presents a longer tail than the Gaussian function. This is the signature of the non-linear structures formed at small scales. Indeed, the PDF of mock galaxy cata-

logs reflects the dynamics that has given rise to structures in the simulations. In this respect, correlations provided by theoretical models and given as initial conditions of the simulations are of CDM type. In these models gravitational clustering forms non-linear structures (i.e. large fluctuations in the density field) up to the limited range of about 10 Mpc/h. Such a length scale can be easily predicted by considering the initial amplitude of fluctuations in these models, normalized to CMBR anisotropies, and by computing the evolution, in an expanding universe, of the linear regime of gravitational clustering. Note that Einasto et al. (2006) reached a similar conclusion, i.e. that in the 2dFGRS there is a number of rich super-clusters which is higher than that found in similar volumes in the Millennium simulation.

Our conclusion is therefore that non-linear structures (i.e. large fluctuations) predicted by models extend to scales which are much smaller than the ones we have found in the 2dFGRS sample and that amplitude of fluctuations on large scales, i.e. $r > 10$ Mpc/h, is large in the real samples and small in the mock galaxy catalogs. We thus conclude that standard models of galaxy formation, with CMBR normalization, are unable to form such large scale structures as the ones present in the observed galaxy distribution.

In addition we have discussed that standard models, beyond the value of the homogeneity scale, predict a second intrinsic length scale represented by the distance-scale at which the two-point correlation function changes sign, i.e. marking the transition from small amplitude fluctuations with weak positive correlations to the regime where there are anti-correlations (and fluctuations of extremely small amplitude). We have pointed out that this length scale is about $r_c \approx 100$ Mpc/h in current models, i.e. it is of slightly larger than the lower limit to the homogeneity scale $\lambda_0 = 70$ Mpc/h we can place by studying the 2dFGRS samples. Forthcoming galaxy sample will allow us to determine whether there will be a transition to super-homogeneity (negative correlations) and at which scale this eventually occurs.

Acknowledgments

We thank Michael Joyce, Andrea Gabrielli, and Luciano Pietronero for useful remarks and discussions. We are grateful to Martin Lopez-Correidoira and Michael Blanton for interesting comments. We also thank an anonymous referee for useful suggestions. We acknowledge the use of the 2dFGRS data available at <http://www.mso.anu.edu.au/2dFGRS/> and of the Millennium run semi-analytic galaxy catalog available at <http://www.mpa-garching.mpg.de/galform/agnpaper/>.

References

Baryshev, Yu., Teerikorpi P., Bulletin Special Astrophysical Observatory, 2006 59, 92

Benoist, C., Maurogordato, S., da Costa, L. N., Cappi, A., Schaeffer, R., 1996, ApJ, 472, 452
 Buswell, G.S., et al., 2004, MNRAS, 354, 991
 Chiaki H. et al., 2003, PASJ, 55, 911
 Colless M., et al., 2001, MNRAS, 328, 1039
 Croton, D.J., et al., 2006, MNRAS, 365, 11
 da Costa, L. et al., 1988, ApJ, 327, 544
 Davis, M., Peebles, P.J.E., 1983, ApJ, 267, 465
 Davis, M. & Huchra, J. 1982, ApJ, 254, 437
 Davis, M. et al., 1988, ApJ, 333, L9
 de Lapparent, V., Geller, M. J., Huchra, J. P., 1986, ApJ, 302, 1
 de Vaucouleurs, G., 1970, Science, 167, 1203
 Eke, V., et al., 2004 MNRAS, 348, 866
 Einasto, J., et al., 2006, A&A, 459, L1
 Falco, E.F., et al., 1999, PASP, 111, 438,
 Frith, W.J., G.S. Buswell, R. Fong, N. Metcalfe, T. Shanks, 2003, MNRAS, 345, 1049
 Gabrielli, A., Sylos Labini, F., 2001, Europhys.Lett., 54, 286
 Gabrielli, A., Joyce, M., Sylos Labini, F., 2002, Phys.Rev., D65, 083523
 Gabrielli A., Sylos Labini F., Joyce M., Pietronero L., 2005 Statistical Physics for Cosmic Structures (Springer Verlag, Berlin)
 Geller, M., Huchra J., 1989, Science 246, 897
 Giovanelli, R., Haynes, M. P., 1993, AJ, 105, 1271
 Gott, J.R. III, et al., 2005, ApJ, 624, 463
 Hawkins E., et al., 2003, MNRAS, 346, 78
 Hogg D.W., Eisenstein D., Blanton M. et al., 2005, ApJ, 624, 54
 Hogg, D.W., Baldry, I.K., Blanton, M.R., Eisenstein, D.J., 2002, eprint arXiv:astro-ph/0210394v1
 Huchra, Davis, Latham and Tonry, 1983, ApJS 52, 89
 Kaiser, N., 1984, ApJ, 284, L9
 Kerscher, M., Schmalzing, J., Buchert, T. and Wagner, H., 1998 A&A, 333, 1
 Kerscher, M. 1999, A&A, 343, 333
 Joyce, M., Sylos Labini, F., 2001, ApJ, 554, L1
 Landy S. D., Szalay A. 1993, ApJ, 412, 64
 Madgwick D. S., et al. 2002, MNRAS, 333, 133
 Norberg, P., et al., 2002, MNRAS, 306, 907
 Norberg E., et al., 2001, MNRAS, 328, 64
 Norberg E., et al., 2002, MNRAS, 332, 827
 Park, C., Vogeley, M. S., Geller, M. J., Huchra, J. P., 1994, ApJ, 431, 569
 Peacock, J.A., 1999, Cosmological physics (Cambridge University Press)
 Peebles P. J. E., 1980, The Large-Scale Structure of the Universe, (Princeton University Press)
 Picard, A., 1991, AJ, 102, 445
 Ratcliffe, A., Shanks, T., Parker, Q.A. & Fong, R., 1998a, MNRAS, 293, 197
 Saslaw, W.C., 2000, The Distribution of the Galaxies, (Cambridge University Press, Cambridge)
 Shectman, S. A., et al., 1996, ApJ, 470, 172
 Spergel, D.N., et al., 2007, ApJS, 170, 377
 Springel, V., et al., 2005, Nature, 435, 629

- Sylos Labini, F., Vasilyev, N.L., Pietronero, L., Baryshev, Yu. V., 2008, eprint arXiv:0805.1132v1
- Sylos Labini, F., Vasilyev, N.L., 2008, A&A, 477, 381
- Sylos Labini, F., Montuori, M., Pietronero, L., 1988, Phys.Rep., 293, 61
- Sylos Labini, F., Vasilyev, N.L. Baryshev, Yu.V., 2007, A&A, 465, 23
- Torquato, S. & Stillinger, F. H., 2003, Phys. Rev., E, 68, 041113
- Totsuji, H., Kihara, T., 1969, PASJ, 21, 221
- Vasilyev, N.L., Baryshev, Yu. V., Sylos Labini, F., 2006, A&A, 447, 431
- York, D., et al., 2000, AJ, 120, 1579
- Weir N., Djorgovski, G.S., Fayyad, U.M., 1995, AJ, 110, 1
- Wu, K.K., Lahav, O. & Rees, M., 1999, Nature, 225, 230
- Zehavi, I., et al., 2002, ApJ, 571, 172
- Zehavi, I., et al., 2004, ApJ, 608, 16
- Zehavi, I., et al., 2005, ApJ, 621, 22